

CLAIMS

What is claimed is:

- Sub A*
1. A method of treating glycogen storage disease type II in an individual, *also form of disease or acid maltase deficiency*
 5 comprising administering to the individual a therapeutically effective amount of human acid α -glucosidase at a regular interval.
2. The method of Claim 1, wherein the glycogen storage disease type II is infantile glycogen storage disease type II.
3. The method of Claim 1, wherein the glycogen storage disease type II is juvenile glycogen storage disease type II.
- 10 4. The method of Claim 1, wherein the glycogen storage disease type II is adult-onset glycogen storage disease type II.
5. The method of Claim 1, wherein the therapeutically effective amount of human acid α -glucosidase is less than about 15 mg of acid α -glucosidase per kilogram of body weight of the individual.
- 15 6. The method of Claim 5, wherein the therapeutically effective amount of human acid α -glucosidase is about 1-10 mg of acid α -glucosidase per kilogram of body weight of the individual.
7. The method of Claim 5, wherein the therapeutically effective amount of human acid α -glucosidase is about 5 mg of acid α -glucosidase per kilogram of body
 20 weight of the individual.

FOOT-20-TSHE000

Sub
A2

8. The method of Claim 1, wherein the human acid α -glucosidase is recombinant human acid α -glucosidase.

9. The method of Claim 1, wherein the human acid α -glucosidase is a precursor of recombinant human acid α -glucosidase.

5 10. The method of Claim 9, wherein the recombinant human acid α -glucosidase is produced in Chinese hamster ovary cells.

11. The method of Claim 1, wherein the regular interval is monthly.

12. The method of Claim 1, wherein the regular interval is bimonthly.

13. The method of Claim 1, wherein the regular interval is weekly.

10 14. The method of Claim 1, wherein the regular interval is twice weekly.

15. The method of Claim 1, wherein the regular interval is daily.

16. The method of Claim 1, wherein the human acid α -glucosidase is administered intravenously.

15 17. The method of Claim 1, wherein the human acid α -glucosidase is administered intramuscularly.

18. The method of Claim 1, wherein the human acid α -glucosidase is administered intrathecally or intraventricularly.

FOOTNOTES

Sub B27

B

